arXiv:hep-ph/0004037v2 7 Apr 2000

About the Mixing and CP Violation in Neutrino System

Yong Liu[1,2] and Utpal Sarkar[3]

1. Laboratory of Numeric Study for Heliospheric Physics
Chinese Academy of Sciences
P. O. Box 8701, Beijing 100080, P.R.China
2. Physics Department, National University of Singapore
Kent Ridge, Singapore 119260
E-mail: phyliuy@nus.edu.sg
3. Physical Research Laboratory, Ahmedabad 380 009, India

Abstract

Suppose the geometrical explaination to the weak CP phase in quark sector is also valid for neutrinos, the mixing and CP violation in neutrino system are discussed. We find a larger J_{CP} than 3×10^{-3} implys the large-mixing solution for solar neutrino problem. In case of bi-maximal mixing, we predict relative large CP violation with J_{CP} larger than 10^{-3} in neutrino system, except the third mixing angle approachs to 0 or $\pi/2$ very closely.

PACS number(s): 11.30.E, 14.60.P, 12.15.F

1. Introduction

In the Standard Model, photon and neutrino have no masses. However, the masslessness of photon is ensured by gauge invariance while the masslessness of neutrinos is only an artificial supposition. Since it is found that, the observed solar neutrino fluxes are all below the predictions based on the Standard Solar Model (SSM) [1], and it is very difficult to explain these flux deficits by modifying the SSM [2], people began to guess that neutrinos may have non-zero masses, and can oscillate from one flavor to another like that occuring in the sector of quarks.

Recently, the Super-Kamiokande experiment has provided a strong evidence for non-zero masses and oscillations of neutrinos [3, 4]. Because it is the first sign for new physics beyond the Standard Model, it has brought up a turbulent shock in the research field of particle physics after the announcement of the Super-Kamiokande result.

In analogy to the quark mixing in the Standard Model, it is expected that, the mixing matrix of the neutrino sector has the similar structure to that of quark sector. Then, it remind us to discuss the problem of mixing and CP violation in neutrino system naturally [5, 6, 7, 8].

Based on the postulation put forward by one of us and the collaborators [9, 10], we investigate the mixing and CP violation in neutrino system in this work. Here, we suppose that

A. The postulation on the relation between weak CP phase and the other three mixing angles in Cabibbo-Kobayashi-Maskawa matrix for quark sector be also available for neutrino system. Due to the similarity between quark sector and lepton sector, we take this supposition as reasonable.

B. Neutrinos come in three families with no additional species, sterile etc., and their masses are hierarchical with m_{ν_e} being smallest and m_{ν_τ} largest [11].

C. Due to the confirmation of the Liquid Scintillator Neutrino Detector (LSND) results at Los Alamos awaiting future experiments [12], in the simplest explanation, solar neutrino data can be understood in terms of $\nu_e - \nu_\mu$ oscillation with a small mass splitting not to influence atmospheric data, and atmospheric data can be explained in terms of $\nu_\mu - \nu_\tau$ large mixing with a large mass splitting compared to the $\nu_e - \nu_\mu$ case [13].

In fact, supposition A is necessary in this paper, while the other two suppositions are only for the convenience. Although we need two mixing angles precisely here, we need not limit which two of the three mixing angles. The suppositions B and C are helpful to draw a clear physical picture.

2. The postulation on the weak CP phase

In previous works [9, 10], we have postulated that, the weak CP phase δ_{13} and the other three mixing angles θ_{12}, θ_{23} and θ_{13} satisfy

$$\sin\delta_{13} = \frac{(1 + s_{12} + s_{23} + s_{13})\sqrt{1 - s_{12}^2 - s_{23}^2 - s_{13}^2 + 2s_{12}s_{23}s_{13}}}{(1 + s_{12})(1 + s_{23})(1 + s_{13})} \tag{1}$$

where, the convention $s_{ij} = \sin\theta_{ij}, c_{ij} = \cos\theta_{ij}$ (the "generation" labels $i, j = 1, 2, 3$) are used and, δ_{13} and θ_{ij} are those present in the standard parametrization of the Cabibbo-Kaboyashi-Maskawa (CKM) matrix

$$V_{KM} = \begin{pmatrix} c_{12}c_{13} & s_{12}c_{13} & s_{13}e^{-i\delta_{13}} \\ -s_{12}c_{23} - c_{12}s_{23}s_{13}e^{i\delta_{13}} & c_{12}c_{23} - s_{12}s_{23}s_{13}e^{i\delta_{13}} & s_{23}c_{13} \\ s_{12}s_{23} - c_{12}c_{23}s_{13}e^{i\delta_{13}} & -c_{12}s_{23} - s_{12}c_{23}s_{13}e^{i\delta_{13}} & c_{23}c_{13} \end{pmatrix} \tag{2}$$

with the real angles θ_{12}, θ_{23} and θ_{13} can all be made to lie in the first quadrant. The phase δ_{13} lies in the range $0 < \delta_{13} < 2\pi$. In following, we will fix the three angles θ_{12}, θ_{23} and θ_{13} in the first quadrant.

The geometry meaning of Eq.(1) is evident. δ_{13} is the solid angle enclosed by $(\pi/2 - \theta_{12}), (\pi/2 - \theta_{23})$ and $(\pi/2 - \theta_{13})$ standing on a same point, or, the area to which the solid angle corresponding on a unit spherical surface.

Hence, to make $(\pi/2 - \theta_{12}), (\pi/2 - \theta_{23})$ and $(\pi/2 - \theta_{13})$ be able to enclose a solid angle, the following relation must be hold.

$$(\frac{\pi}{2} - \theta_{ij}) + (\frac{\pi}{2} - \theta_{jk}) \geq (\frac{\pi}{2} - \theta_{ki}) \qquad (i \neq j \neq k \neq i = 1, 2, 3. \quad \theta_{ij} = \theta_{ji}) \tag{3}$$

Eq.(3) and Eq.(1) are the most important constraints in this work, on which the following discussions are based.

3. The relevant experimental results on neutrino masses and mixing

The recent analysis made by Hata and Langacker [14] gives viable solutions for the BAHCALL SSM [15, 16]. With the Mikheyev-Smirnov-Wolfenstein (MSW) mechanism being considered [17, 18], they give the small-mixing solution

$$\delta m_{sol}^2 \sim 5 \times 10^{-6} eV^2 \qquad sin^2 2\theta_{sol} \sim 8 \times 10^{-3} \tag{4}$$

and the large-mixing solution

$$\delta m_{sol}^2 \sim 1.6 \times 10^{-5} eV^2 \qquad sin^2 2\theta_{sol} \sim 0.6. \tag{5}$$

Vacuum oscillation also provide solutions

$$\delta m_{sol}^2 = (5 \sim 8) \times 10^{-11} eV^2 \qquad sin^2 2\theta_{sol} = 0.65 \sim 1. \tag{6}$$

The atmospheric neutrino data from Super-Kamiokande etc. imply that the parameters of the $\nu_\mu - \nu_\tau$ oscillation of the atmospheric neutrinos are [3, 4, 16]

$$10^{-4} \leq \delta m^2_{atm} \leq 10^{-2} eV^2 \qquad sin^2 2\theta_{atm} \approx 1. \tag{7}$$

We have known that, the small-mixing solution causes the energy-spectrum distortion while the large-mixing solution causes the day-night flux difference, and, the vacuum-oscillations cause seasonal variation of the ^7B_e solar neutrino flux [16].

In the next section, we will see that, with the Super-Kamiokande results about the $\nu_\mu - \nu_\tau$ oscillation being admitted, the small-mixing solution differents from the large-mixing solution in CP violation greatly. Because the discussions here are only related to the mixing angles, and the MSW large-mixing solution gives about the same mixing as that given by the vacuum oscillation solution, so, we only discuss the two cases of small- and large-mixing as indicated by the MSW solutions.

4. Some predictions on the mixing and CP violation in neutrino system

Let us return to the constraint Eq.(3) and recall the clear geometry meaning of Eq.(1), what can we extract from them? As we have supposed in section 1, Eq.(7) tells us

$$\theta_{\mu\tau} \approx \pi/4. \tag{8}$$

Keep this point in mind, then, by the use of Eq.(3), Eq.(4-6) will give restriction on the mixing angle between ν_e and ν_τ. In the meantime, Eq.(1) tell us some information on the CP violation in neutrino system. Let us talk more detail about these two problems.

(1) About the mixing angle between ν_e and ν_τ.

From Eq.(3), we have

$$|(\frac{\pi}{2} - \theta_{e\mu}) - (\frac{\pi}{2} - \theta_{\mu\tau})| \leq (\frac{\pi}{2} - \theta_{e\tau}) \leq Min(\pi/2, \ (\frac{\pi}{2} - \theta_{e\mu}) + (\frac{\pi}{2} - \theta_{\mu\tau})) \tag{9}$$

Note that, Eqs.(4-5) imply for small- and large-mixing solutions

$$\theta_{e\mu} \sim 0.045 \quad \text{or} \quad \pi/2 - 0.045$$

and

$$\theta_{e\mu} \sim 0.443 \quad \text{or} \quad \pi/2 - 0.443$$

respectively. Considering Eq.(8), then we obtain

$$0 \leq \theta_{e\tau} \leq \pi/4 + 0.045 \tag{10}$$

or

$$\pi/4 - 0.045 \leq \theta_{e\tau} \leq \pi/4 + 0.045 \tag{11}$$

for the case of small-mixing solution. And

$$0 \le \theta_{e\tau} \le \pi/4 + 0.443 \tag{12}$$

or

$$\pi/4 - 0.443 \le \theta_{e\tau} \le \pi/4 + 0.443 \tag{13}$$

for the case of large-mixing solution. Eq.(10-13) are the constraints on the mixing angle $\theta_{e\tau}$.

(2) About the CP violation in neutrino system.

Now, we discuss the CP violation in neutrino system. As is well known, all the CP violated observables are proportional to the Jarlskog invariant [19, 20]

$$J_{CP} = s_{12}s_{13}s_{23}c_{12}c_{13}^2c_{23}s_{\delta_{13}} \tag{14}$$

where, $s_{\delta_{13}} \equiv sin\delta_{13}$ with δ_{13} the weak CP phase presenting in the CKM matrix of the lepton sector. And, we instead θ_{12}, θ_{13} and θ_{23} by $\theta_{e\mu}, \theta_{e\tau}$ and $\theta_{\mu\tau}$ respectively in following.

With $\theta_{\mu\tau}$ given by the Super-Kamiokande data definately, the small- and large-mixing solutions and the vacuum oscillation will give the pemitted ranges for $\theta_{e\tau}$ correspondingly. Substitute Eq.(1) into Eq.(14), we obtain J_{CP} as a function of $\theta_{e\tau}$. Then, we can draw the curve which J_{CP} versus $\theta_{e\tau}$. The results are shown in Fig.(1). From the figure, we find that, the small-mixing solution corresponds to small CP violation while the large-mixing solution corresponds to large CP violation.

For the small-mixing solution, J_{CP} is very small. The maximum of J_{CP} is about 5×10^{-3} when $\theta_{e\mu}$ nearing to 0, and 1.5×10^{-4} when $\theta_{e\mu}$ nearing to $\pi/2$.

For the large-mixing solution, J_{CP} is relative large. The maximum of J_{CP} is about 1.5×10^{-2} when $\theta_{e\mu}$ nearing to 0.443, and 3.2×10^{-2} when $\theta_{e\mu}$ nearing to $\pi/2 - 0.443$.

Now, it is evident that, if the future experiments on the CP violation in neutrino system tell us that J_{CP} is larger than 5×10^{-3}, the mixing between ν_e and ν_μ must be large. Recalling the same observable in quark sector obtained via $K^0 - \overline{K^0}$ system is about 10^{-4}, if it is the same order of magnitude in the neutrino system, then, the mixing between ν_e and ν_τ is either around 0.8 with a relative narrow window or nearing to 0 or $(\pi/4 + 0.443)$ very closely.

On the other hand, either $\theta_{e\mu}$ takes ~ 0.443 or it takes $\sim (\pi/2 - 0.443)$ for large-mixing solution can also be distingushed to some extent. For example, if experiment tells us $J_{CP} > 0.015$, it must be $\theta_{e\mu} \sim 0.443$.

Finally, with the CHOOZ result [21] being considered, that is, $\sin^2 2\theta_{e\tau} < 0.2$, then we have

$$0 < \theta_{e\tau} < 0.23 \quad \text{or} \quad \pi/2 - 0.23 < \theta_{e\tau} < \pi/2.$$

Based on the above constraint, we can see from Fig.(1), firstly, the small-mixing solution with $\theta_{e\mu} \sim (\pi/2 - 0.045)$ and the large-mixing solution with $\theta_{e\mu} \sim (\pi/2 - 0.443)$ have been excluded.

Secondly, the possible domain $\pi/2 - 0.23 < \theta_{e\tau} < \pi/2$ should also be eliminated. Thirdly, J_{CP} can still be large to about 0.02 in the case of large-mixing solution with $\theta_{e\mu} \sim 0.443$, especially, the larger J_{CP} than 3×10^{-3} will exclude the possibility of small-mixing solution finally.

Maybe, the most interesting conclusion is about the bi-mixmal mixing [22, 23]. From the above analysis, we can see that, in most of the permitted range of the third angle - $\theta_{e\tau}$, there will be a relative large CP violation.

Suppose that $\theta_{e\mu} = \theta_{\mu\tau} = \pi/4$, similarly, we can draw the curve which J_{CP} versus $\theta_{e\tau}$ in the permitted range of $\theta_{e\tau}$. The result is shown in Fig.(2). We find that, except for $\theta_{e\tau}$ nearing to 0 or $\pi/2$, J_{CP} is larger than 10^{-3} in most of the permitted range of $\theta_{e\tau}$. And, the maximum of J_{CP} can reach to about 0.018 when the CHOOZ result is considered.

5. Conclusions

Starting from the postulation on the relation between weak CP phase and the other three mixing angles in the CKM matrix, we have investigated the mixing and CP violation in the neutrino system.

We suppose that, the solar neutrino problem be understood in terms of $\nu_e - \nu_\mu$ oscillation with a small mass splitting. With the definite large mixing between $\nu_\mu - \nu_\tau$ indicated by the Super-Kamiokande data, and the CHOOZ result being considered, we obtain the relevant constraints on the mixing between ν_e and ν_τ. We find, $0 \leq \theta_{e\tau} \leq 0.23$ is permitted by the small- and the large-mixing solutions.

Besides, the mixing between ν_e and ν_μ is limited as $\theta_{e\mu} \sim 0.045$ for the small-mixing solution or $\theta_{e\mu} \sim 0.443$ for the large-mixing solution. And, a larger J_{CP} than 3×10^{-3} will finally exclude the possibility of small-mixing solution.

Furthermore, if the suppositions B and C in section **1** holds, a J_{CP} larger than 3×10^{-3} implys the large-mixing solution for solar neutrino problem. And, if it takes the same order for J_{CP} in the neutrino system as the one in quark system, the mixing between ν_e and ν_μ will be very small ($\sim 10^{-2}$ or less).

For the case of bi-maximal mixing, we predict a large CP violation in neutrino system with J_{CP} larger than 10^{-3}, except the third mixing angle approachs to 0 or $\pi/2$ very closely.

Finally, although we have made some suppositions in this work, the basis and the method used here is actually valid for a more general discussion.

Notes: In fact, this work has been finished and submitted before the last July. Due to some reason, we have not put it on the net in time. Just two days ago, when we noted the paper hep-ph/0004020 by Sin Kyu Kang, C. S. Kim and J. D. Kim and found their results are almost the same as those of us, although they based on some concrete model while we only started out from our postulation, we are encouraged to post this short paper.

References

[1] J. Bahcall, S. Basu and M. Pinsonneault, Phys. Lett. **B** 433, 1(1998).

[2] J. Bahcall and H. Bethe, Phys. Rev. Lett. **65**, 2233(1990) and Phys. Rev. D**44**, 2962(1991). N. Hata and P. Langacker, Phys. Rev. D**56**, 6107(1997).

[3] Super-Kamiokande Collaboration : Y. Fukuda *et al*, Phys. Rev. Lett. **81**, 1562 (1998); hep-ex/9805006; Phys. Lett. **B 433**, 9 (1998); T. Kajita, hep-ex/9810001.

[4] Super-Kamiokande Collaboration: Y. Fukuda *et al*, Phys. Rev. Lett. **81**, 1158 (1998); Talk by Y. Suzuki at Neutrino'98, Takayama, Japan (1998).

[5] Utpal Sarkar and R. Vaidya, Phys. Lett. **B 442**, 243(1998). C. S. Kim, J. D. Kim, Phys. Rev. **D 61**, 057302(2000). Sin Kyu Kang, C. S. Kim and J. D. Kim, hep-ph/0004020. A. Mondragon and E. Rodri Guez-Jourgui, Phys. Rev. **D 59**, 093009(1999).

[6] K. R. Schubert, May we expect CP- and T-violation effects in neutrino oscillation ? hep-ph/9902215. De Rujula, M. B. Gavela and P. Hernandez, hep-ph/9811390, Nucl. Phys. **B 547**, 21(1999). A. Donini, M. B. Gavela, P. Hernandez and S. Rigolin, hep-ph/9909254. A. Cervera, A. Donini, M. B. Gavela, J. J. Gomez Cadenas, P. Hernandez, O. Mena and S. Rigolin, hep-ph/0002108.

[7] A. M. Gago, V. Pleitez and R. Zukanovich Funchal, CP violation in vacuum neutrino oscillation experiments, hep-ph/9810505.

[8] V. Barger, K. Whisnant and R. J. N. Phillips, Phys. Rev. Lett. **45**, 2084(1980).

[9] Jing-Ling Chen, Mo-Lin Ge, Xue-Qian Li and Yong Liu, A Possible Hidden Symmetry and Geometrical Source of The Phase in The CKM Matrix. Eur. Phys. J. C **9**, 437(1999).

[10] Yong Liu, hep-ph/9910234. Phys. Rev. **D 61**, 033010(2000).

[11] P. M. Fishbane and P. Kaus, On neutrino masses and leptonic mixing, hep-ph/9810233.

[12] C. Athanassopoulos et al. (LSND Collaboration), Phys. Rev. Lett. **75**, 2650(1995); ibid. **77**, 3082(1996).

[13] R. Barbieri et al., hep-ph/9807235. E. Malkawi, Common hierarchical lepton mass textures for atmospheric and solar neutrino oscillations, hep-ph/9810542.

[14] N. Hata and P. Langacger, Phys. Rev. D**56**, 6107(1997).

[15] J. N. Bahcall and M. H. Pinsonneault, Rev. Mod. Phys. **67**, 781(1995).

[16] Particle Data Group, The Europ. Phys. J. C**3**, 310 -329(1998).

[17] L.Wolfenstein, Phys.Rev.D 17(1978)2369; 20(1979)2634.

[18] S.P.Mikheyev and A.Yu.Smirnov, Yad.Fiz. 42(1985)1441(Sov.J.Nucl.Phys.42(1985)913); Nuov.Cimento.C 17(1986)9.

[19] *CP Violation* Ed. C. Jarlskog, World Scientific Publishing Co. Pte. Ltd 1989.

[20] E. A. Paschos and U. Turke, Phys. Rept.**4**, 145(1989).

[21] CHOOZ Collaboration : M. Appollonio et al, hep-ex/9711002.

[22] V. Barger, S. Pakvasa, T. J. Weiler and K. Whisant, Phys. Lett. **B 437**, 107(1998). C. Giunti, hep-ph/9810272. D. V. Ahluwalia, Mod. Phys. Lett. **A 13**, 2249(1998).

[23] M. Tanimoto, Phys. Rev. D**59**, 017304(1999). I. Stancu and D. V. Ahluwalia, Phys. Lett. **B 460**, 431(1999).



Figure 1: J_{CP} versus $\theta_{e\tau}$. Where, $\theta_{\mu\tau} = \pi/4$. The curves s1, s2, l1 and l2 corresponds to the cases of $\theta_{e\mu} = 0.045,\ (\pi/2 - 0.045),\ 0.443$ and $(\pi/2 - 0.443)$ respectively.



Figure 2: J_{CP} versus $\theta_{e\tau}$. Where, $\theta_{e\mu} = \theta_{\mu\tau} = \pi/4$. So, the permitted range for $\theta_{e\tau}$ is $0 \sim \pi/2$.